

An analog of the vaisman-molino cohomology for manifolds modelled on some types of modules over weil algebras and its application

Shurygin V., Smolyakova L.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

An epimorphism $\mu : A \rightarrow B$ of local Weil algebras induces the functor T_μ from the category of fibered manifolds to itself which assigns to a fibered manifold $p : M \rightarrow N$ the fibered product $p_\mu : T_A N \times_{T_B N} T_B M \rightarrow T_A N$. In this paper we show that the manifold $T_A N \times_{T_B N} T_B M$ can be naturally endowed with a structure of an A -smooth manifold modelled on the A -module $L = A^n \oplus B^m$, where $n = \dim N$, $n + m = \dim M$. We extend the functor T_μ to the category of foliated manifolds (M, F) . Then we study A -smooth manifolds ML whose foliated structure is locally equivalent to that of $T_A N \times_{T_B N} T_B M$. For such manifolds ML we construct bigraduated cohomology groups which are similar to the bigraduated cohomology groups of foliated manifolds and generalize the bigraduated cohomology groups of A -smooth manifolds modelled on A -modules of the type A^n . As an application, we express the obstructions for existence of an A -smooth linear connection on ML in terms of the introduced cohomology groups.
